

The Schrodinger Equation From a Quadratic Hamiltonian System

Wai Bong Yeung*

Institute of Physics, Academia Sinica, Taipei, Taiwan, ROC

(Oct 1999)

We regard the real and imaginary parts of the Schrodinger wave function as canonical conjugate variables. With this pair of conjugate variables and some other $2n$ pairs, we construct a quadratic Hamiltonian density. We then show that the Schrodinger Equation follows from the Hamilton's Equation of motion when the Planck frequency is much larger than the characteristic frequencies of the Hamiltonian system. The Hamiltonian and the normal mode solutions coincide, respectively, with the energy expectation value and the energy eigenstates of the corresponding quantum mechanical system.

Let us consider a Hamiltonian system consists of $(1+2n)$ pairs of canonical conjugate variables (p, q) (P_j, Q_j), (π_j, η_j) , $j=1, \dots, n$. All of these canonical variables are function of $x = (x_1, \dots, x_n)$ and t .

The Hamiltonian density $H(p, q; P_j, Q_j; \pi_j, \eta_j; \partial q, \partial Q, \partial \eta)$ is quadratic in the variables and chosen to be

$$H = 1/2(V(x)(p^2 + q^2) - m(P_j^2 + Q_j^2 + \pi_j^2 + \eta_j^2) - p\partial_j(Q_j + \eta_j) - (P_j + \pi_j)\partial_j q) \quad (1)$$

where m is a given parameter of the system. The function $V(x)$ is also given. ∂_j denotes the differentiation with respect of x_j . Repeated indices will be summed over from 1 to n .

The $(2+4n)$ Hamilton Equation of motion, obtained by varying independently the canonical variables [1] are

$$\begin{aligned} \partial_t p &= -V(x)q - 1/2\partial_j(P_j + \pi_j) \\ \partial_t q &= V(x)p - 1/2\partial_j(Q_j + \eta_j) \end{aligned} \quad (2)$$

Similarly

$$\begin{aligned} \partial_t P_i &= mQ_i - 1/2\partial_i p \\ \partial_t Q_i &= -mP_i - 1/2\partial_i q \\ \partial_t \pi_i &= m\eta_i - 1/2\partial_i p \\ \partial_t \eta_i &= -m\pi_i - 1/2\partial_i q \end{aligned} \quad (3)$$

These are coupled linear equations, and hence linear superpositions of solutions of these equations are also solutions to these equations. In other words, the principle of linear superposition [2] applies in this Hamiltonian system. Now consider the case in which m is a very large parameter and that (P_i, Q_i) and (π_i, η_i) are slowly varying variables. In that case Eq(3) can be approximated by

$$\begin{aligned} Q_i &= 1/2m\partial_i p \\ P_i &= -1/2m\partial_i q \\ \eta_i &= 1/2m\partial_i p \\ \pi_i &= -1/2m\partial_i q, j = 1, \dots, n \end{aligned} \quad (4)$$

And because m is a very large number, (P_i, Q_i) and (π_i, η_i) are small relative to (p, q) . Eq(4) gives the relations between the large variables (p, q) and the small variables (P_i, Q_i) , (π_i, η_i) . Eliminating the small variables from Eq(2), we get the following equations containing the large variables only.

$$\begin{aligned} \partial_t p &= -V(x)q + 1/2m\partial^2 q \\ \partial_t q &= V(x)p - 1/2m\partial^2 p \end{aligned} \quad (5)$$

Let us now construct a complex function $\psi(x, t)$ by using the above $q(x, t)$ and $p(x, t)$ as the real and imaginary parts of $\psi(x, t) = (q(x, t) + ip(x, t))/\sqrt{2}$. It can be seen immediately that Eq(5) can be summarized as the real and imaginary parts, respectively, of the complex equation

*Electronic address: phweung @ccvax.sinica.edu.tw

$$i\partial_t\psi = (-1/2m\partial^2 + V(x))\psi \quad (6)$$

This complex equation is nothing but the Schrodinger equation in natural units ($\hbar=c=1$). The parameter m and the function $V(x)$ will then be, respectively, the mass and the potential energy of the corresponding quantum mechanical system.

It is interesting to calculate the Hamiltonian $H = \int d^n x H$ for the system. With the relations given in Eq(4) substituted into Eq(1), the Hamiltonian of the system will read as

$$H = \int d^n x 1/2(V(x)(p^2 + q^2) - m(1/2m^2(\partial_j q)^2 + 1/2m^2(\partial_j p)^2) - 1/mp\partial^2 p + 1/m(\partial_j q)^2)) \quad (7)$$

Integration by parts and discarding surface terms, we get

$$H = \int d^n x 1/2(V(x)(p^2 + q^2) - p1/2m\partial^2 p - q1/2m\partial^2 q) \quad (8)$$

Which is the same as $\int d^n x \psi * (-1/2m\partial^2 + V(x))\psi$. And hence the Hamiltonian H of our Hamiltonian system is just the expectation value of the nonrelativistic energy of the corresponding wave function.

Furthermore ,Eq(5) can be used to rewrite the Hamiltonian H into the the form

$$H = \int d^n x 1/2(p\partial_t q - q\partial_t p) \quad (9)$$

This H could be a function of time even if $p(x, t)$ and $q(x, t)$ satisfy Eq(5) If we are interested in looking for normal mode solutions for the Hamiltonian system , then we should look for solutions which will give us an effective Hamiltonian of the form of

$$H = E \int d^n x 1/2(p^2 + q^2) \quad (10)$$

Where E is a constant to be determined . Comparision of Eq(9)with Eq(10) will give

$$\begin{aligned} \partial_t q &= Ep \\ \partial_t p &= -Eq \end{aligned} \quad (11)$$

Eq(11) will ensure that H be time independent and will also cast Eq(5) into the form of

$$\begin{aligned} Eq &= V(x)q - 1/2m\partial^2 q \\ Ep &= V(x)p - 1/2m\partial^2 p \end{aligned} \quad (12)$$

This Eq(12) is exactly the same as the eigenvalue problem for

$$E\psi = (V(x) - 1/2m\partial^2)\psi \quad (13)$$

To consider the Hamiltonian system in MKS units, we have to put in some dimensional parameters to take care of the different dimensionalities of various terms of the Hamiltonian density. If we choose p and q to be dimensionless , then the Hamilton's Equation of motion will require H to be 1/time in dimension . And since $V(x)$ is the potential energy, the first term on the right hand side of Eq(10) must be divided by a constant of the dimension of energy-time in order to get the dimension right. This unique constant is, of course, the Planck's constant $\hbar$. Simple arguments will fix the other constants before variuos terms , and the result is the following Hamiltonian density

$$H = (1/2\hbar)(V(x)(p^2 + q^2) - (1/2\hbar)(mc^2)(P_j^2 + Q_j^2 + \pi_j^2 + \eta_j^2) - (c/2)p\partial_j(Q_j + \eta_j) - (c/2)(P_j + \pi_j)\partial_j q) \quad (14)$$

This Hamiltonian will generate the Schrodinger Equation

$$i\hbar\partial_t\psi = (-\hbar^2/2m\partial^2 + V(x))\psi \quad (15)$$

under the conditions that we have mentioned above. And Eq(15), which in turn, will imply the prescription $E \rightarrow i\hbar\partial_t$, momemtum $\rightarrow -i\hbar\partial_x$, used in quantum mechanics for $E = (momentum)^2/2m + V(x)$.

The very large parameter will now be the Planck frequency $mc^2/\hbar$. The low frequency sector of the Hamiltonian system , which is the sector with characteristic freqeicies considerably smaller than the Planck frequency , will coincide

exactly with the Schrodinger quantum mechanics. Our restriction to the low frequency sector of the Hamiltonian system , we think , is not a drawback of our approach because of the fact that the Schrodinger quantum mechanics is a nonrelativistic limit by itself and also that the Planck frequency is always a gigantic number for all known atomic particle In this approach of obtaining the Schrodinger Equation, two more things worth our attention. Firstly, the Planck constant h appears as a parameter in the Hamiltonian density. The role thus played by h in our approach is radically different from that played in the conventional presentation of quantum mechanics. Secondly ,physically there are the small hidden variable. They come from the negative parts of the Hamiltonian density, and are playing important roles in the determination of the evolution of $p(x, t)$ and $q(x, t)$, and hence $\psi(x, t)$ [3] .

- [1] .Goldstein, Classical Mechanics,Addison-Wesley Press. Cambridge.Ma 1980
- [2] .A.M.Dirac ,Quantum Mechanics.Oxford University Press,New York.1974.
- [3] .Bohm, Quantum Mechanics,Prentice Hall, New York.1951